Mathematische Annalen manuscript No.
(will be inserted by the editor)

On the distribution of imaginary parts of zeros of the Riemann zeta function, II

Kevin Ford ⋆ · K. Soundararajan ⋆⋆ · Alexandru Zaharescu ⋆⋆⋆

the date of receipt and acceptance should be inserted later – © Springer-Verlag 2019

Abstract. *Mathematics Subject Classification (2000):* Primary 11M26; Secondary 11K38 We continue our investigation of the distribution of the fractional parts of $\alpha\gamma$, where α is a fixed non-zero real number and γ runs over the imaginary parts of the non-trivial zeros of the Riemann zeta function. We establish some connections to Montgomery's pair correlation function and the distribution of primes in short intervals. We also discuss analogous results for a more general L-function.

Key words. Riemann zeta function – zeros, fractional parts – primes in short intervals – pair correlation functions

arXiv:0805.2745v2 [math.NT] 22 Aug 2008

1. Introduction and Statement of Results

In this paper we continue the study of the distribution of the fractional parts $\{\alpha\gamma\}$ initiated by the first and third authors in [3], where α is a fixed positive real number and γ runs over the positive ordinates of zeros of the Riemann zeta function $\zeta(s)$. We extend and generalize the results from [3] in several directions, establishing connections between these fractional parts, the pair correlation of zeros of $\zeta(s)$ and the distribution of primes in short intervals. It is known [9] that for any fixed α, the fractional parts $\{\alpha\gamma\}$ are uniformly distributed (mod 1). That is, for all continuous

Kevin Ford and Alexandru Zaharescu
Department of Mathematics, 1409 West Green Street, University of Illinois at Urbana-Champaign, Urbana, IL 61801, USA

K. Soundararajan
Department of Mathematics, 450 Serra Mall, Bldg. 380, Stanford University, Stanford, CA 94305, USA

⋆ The first author is supported by National Science Foundation Grant DMS-0555367

⋆⋆ The second author is partially supported by the National Science Foundation and the American Institute of Mathematics (AIM)

⋆⋆⋆ The third author is supported by National Science Foundation Grant DMS-0456615

functions $f : \mathbb{T} \to \mathbb{C}$, as $T \to \infty$ we have

$$\sum_{0<\gamma\le T} f(\alpha\gamma) = N(T)\int_{\mathbb{T}} f(x)dx + o(N(T)). \tag{1.1}$$

Here $\mathbb{T}$ is the torus $\mathbb{R}/\mathbb{Z}$ and $N(T)$ denotes the number of ordinates $0 < \gamma \le T$; it is well-known that

$$N(T) = \frac{T}{2\pi}\log\frac{T}{2\pi e} + O(\log T). \tag{1.2}$$

We are interested in the lower order terms in the asymptotic (1.1). For a general continuous function f the asymptotic (1.1) can be attained arbitrarily slowly so that no improvement of the error term there is possible. But if we assume that f has nice smoothness properties then we can isolate a second main term of size about T. More precisely, we define the function $g_\alpha : \mathbb{T} \to \mathbb{C}$ as follows. If α is not a rational multiple of $\frac{\log p}{2\pi}$ for some prime p, then g_α is identically zero. If $\alpha = \frac{a}{q}\frac{\log p}{2\pi}$ for some rational number a/q with $(a, q) = 1$ then we set

$$g_\alpha(x) = -\frac{\log p}{\pi}\Re\sum_{k=1}^{\infty}\frac{e^{-2\pi iqkx}}{p^{ak/2}} = -\frac{(p^{a/2}\cos 2\pi qx - 1)\log p}{\pi(p^a - 2p^{a/2}\cos 2\pi qx + 1)}. \tag{1.3}$$

Then, we expect (for suitable f) that as $T \to \infty$

$$\sum_{0<\gamma\le T} f(\alpha\gamma) = N(T)\int_{\mathbb{T}} f(x)dx + T\int_{\mathbb{T}} f(x)g_\alpha(x)dx + o(T). \tag{1.4}$$

As remarked above, certainly (1.4) does not hold for all continuous functions f. In Corollary 2 of [3], it is shown that (1.4) holds for all $f \in C^2(\mathbb{T})$, and if the Riemann Hypothesis (RH) is true then (1.4) holds for all absolutely continuous functions f (see Corollary 5 there). Moreover it is conjectured there (see Conjecture A there) that (1.4) does hold when f is the characteristic function of an interval in $\mathbb{T}$.

Conjecture 1. Let $\mathbb{I}$ be an interval of $\mathbb{T}$. Then

$$\sum_{\substack{0<\gamma\le T\\ \{\alpha\gamma\}\in\mathbb{I}}} 1 = |\mathbb{I}|N(T) + T\int_{\mathbb{I}} g_\alpha(x)dx + o(T).$$

We define the discrepancy of the sequence $\{\alpha\gamma\}$ (for $0 < \gamma \le T$) as

$$D_\alpha(T) = \sup_{\mathbb{I}}\Big|\frac{1}{N(T)}\sum_{\substack{0<\gamma\le T\\ \{\alpha\gamma\}\in\mathbb{I}}} 1 - |\mathbb{I}|\Big|,$$

where the supremum is over all intervals $\mathbb{I}$ of $\mathbb{T}$. Unconditionally, Fujii [4] proved that $D_\alpha(T) \ll \frac{\log\log T}{\log T}$ for every α. On RH, Hlawka [9] showed that $D_\alpha(T) \ll \frac{1}{\log T}$, which is best possible for α of the form $\frac{a}{q}\frac{\log p}{2\pi}$ ([3], Corollary 3). Conjecture 1 clearly implies the following conjecture for the discrepancy (see Conjecture A and Corollary 6 of [3]).

Conjecture 2. We have

$$D_\alpha(T) = \frac{T}{N(T)} \sup_{\mathbb{I}} \Big| \int_{\mathbb{I}} g_\alpha(x)dx \Big| + o\Big(\frac{1}{\log T}\Big).$$

Even assuming RH, we are unable to establish Conjectures 1 and 2. We show here some weaker results towards these conjectures, and how these conjectures would follow from certain natural assumptions on the zeros of $\zeta(s)$, or the distribution of prime numbers.

Theorem 1. *(i) We have unconditionally*

$$D_\alpha(T) \geq \frac{T}{N(T)} \sup_{\mathbb{I}} \Big| \int_{\mathbb{I}} g_\alpha(x)dx \Big| + o\Big(\frac{1}{\log T}\Big).$$

(ii) Assuming RH, for any interval $\mathbb{I}$ of $\mathbb{T}$ we have

$$\Big| \sum_{\substack{0<\gamma\leq T \\ \{\alpha\gamma\}\in\mathbb{I}}} 1 - |\mathbb{I}|N(T) - T\int_{\mathbb{I}} g_\alpha(x)dx \Big| \leq \frac{\alpha}{2}T + o(T).$$

The left side of (1.1) depends strongly on the behavior of the sums $\sum_{0<\gamma\leq T} x^{i\gamma}$.

Conjecture 3. Let $A > 1$ be a fixed real number. Uniformly for all $\frac{T^2}{(\log T)^5} \leq x \leq T^A$ we have

$$\sum_{0<\gamma\leq T} x^{i\gamma} = o(T). \tag{1.5}$$

Theorem 2. *Assume RH. Then Conjecture 3 implies Conjectures 1 and 2.*

Remarks. Assuming RH, (1.5) holds for $x \to \infty$ and $x = o(T^2/\log^4 T)$ as $T \to \infty$ by uniform versions of Landau's formula for $\sum_{0<\gamma\leq T} x^\rho$ [12]. For example, Lemma 1 of [3] implies, for $x > 1$ and $T \geq 2$, that (unconditionally)

$$\sum_{0<\gamma\leq T} x^\rho = -\frac{\Lambda(n_x)}{2\pi}\frac{e^{iT\log(x/n_x)}-1}{i\log(x/n_x)} + O\left(x\log^2(Tx) + \frac{\log T}{\log x}\right), \tag{1.6}$$

where n_x is the nearest prime power to x, and the main term is to be interpreted as $-T\frac{\Lambda(x)}{2\pi}$ if $x = n_x$. This main term is always $\ll T\log x$. On RH, divide both sides of (1.6) by $x^{1/2}$ to obtain (1.5). Unconditionally, one can use Selberg's zero-density estimate to deduce

$$\Big|\sum_{0<\gamma\le T}(x^{i\gamma}-x^{\rho-1/2})\Big| \ll \frac{T\log^2(2x)}{\log T};$$

see e.g. (3.8) of [3]. This gives (1.5) when $\log x = o(\sqrt{\log T})$.

We next relate Conjecture 3 to the distribution of primes in short intervals.

Conjecture 4. For any $\varepsilon > 0$, if x is large and $y \le x^{1-\epsilon}$, then

$$\psi(x+y)-\psi(x) = y + o(x^{\frac12}/\log\log x).$$

Theorem 3. *Assume RH. Conjecture 4 implies Conjecture 3, and hence Conjectures 1 and 2. Conversely, if RH and Conjecture 3 holds, then for all fixed $\varepsilon > 0$, large x and $y \le x^{1-\varepsilon}$,*

$$\psi(x+y)-\psi(x) = y + o(x^{\frac12}\log x).$$

Remarks. Whereas the behavior of the left side of (1.6) is governed by a single prime when x is small, for larger x the sum is governed by the primes in an interval. It has been conjectured ([16], Conjecture 2) that for $x^\varepsilon \le h \le x^{1-\varepsilon}$, $\psi(x+h)-\psi(x)-h$ is normally distributed with mean 0 and variance $h\log(x/h)$. Thus, it is reasonable to conjecture that for every $\varepsilon > 0$,

$$\psi(x+y)-\psi(x)-y \ll_\varepsilon y^{1/2}x^\varepsilon \qquad (1\le y\le x), \tag{1.7}$$

a far stronger assertion than Conjecture 4. It is known that RH implies $\psi(x) = x + O(x^{1/2}\log^2 x)$ (von Koch, 1900).

A statement similar to the second part of Theorem 3 has been given by Gonek ([7], Theorem 4). Assuming RH, Gonek showed that if

$$\sum_{0<\gamma\le T} x^{i\gamma} \ll_\varepsilon Tx^{-1/2+\varepsilon} + T^{1/2}x^\varepsilon$$

holds uniformly for all $x, T \ge 2$ and for each fixed $\varepsilon > 0$, then (1.7) follows.

We also want to describe how to bound the sum $\sum_{0<\gamma\le T} x^{i\gamma}$ in terms of the pair correlation function

$$\mathcal{F}(x,T) = \sum_{0<\gamma,\gamma'\le T} \frac{4x^{i(\gamma-\gamma')}}{4+(\gamma-\gamma')^2}. \tag{1.8}$$

Such bounds have been given by Gallagher and Mueller [5], Heath-Brown [8], and Goldston and Heath-Brown [6]. First we state a strong version of the Pair Correlation Conjecture for $\zeta(s)$.

Conjecture 5. Fix a real number $A > 1$. Uniformly for all $\frac{T^2}{(\log T)^6} \le x \le T^A$ we have

$$\mathcal{F}(x,T) = N(T) + o\left(\frac{T}{\log T}\right) \qquad (T \to \infty).$$

Theorem 4. *Assume RH. Then Conjecture 5 implies Conjecture 3, and therefore also Conjectures 1 and 2.*

Remarks. The original pair correlation conjecture of Montgomery [14] states that

$$\mathcal{F}(x,T) \sim N(T) \qquad (T \to \infty)$$

uniformly for $T \le x \le T^A$, where A is any fixed real number. Tsz Ho Chan [1] has made an even stronger conjecture than Conjecture 5, namely he conjectured that for any $\epsilon > 0$ and any large $A > 1$,

$$\mathcal{F}(x,T) = N(T) + O\left(T^{1-\epsilon_1}\right)$$

if $T^{1+\epsilon} \le x \le T^A$, where $\epsilon_1 > 0$ may depend on ϵ, and the implicit constant may depend on ϵ and A.

In the next section, we prove Theorems 1–4. In section 3 we discuss analogous results for general L-functions.

2. Proof of Theorems 1–4

Proof (Proof of Theorem 1 (i)). Let $\mathbb{I}$ denote an interval of $\mathbb{T}$ for which $|\int_{\mathbb{I}} g_\alpha(x)dx|$ attains its maximum. Let ϵ be a small positive number, and let $h_\epsilon : \mathbb{T} \to \mathbb{R}$ be a smooth function satisfying $h_\epsilon(x) \ge 0$ for all x, $h_\epsilon(x) = 0$ for $\epsilon < x \le 1$, and $\int_{\mathbb{T}} h_\epsilon(x)dx = 1$. Set $f(x) = \int_{\mathbb{T}} h_\epsilon(y)\chi_{\mathbb{I}}(x-y)dy$, where $\chi_{\mathbb{I}}$ denotes the characteristic function of the interval $\mathbb{I}$. Then f is smooth, and so (1.4) holds for f. Therefore

$$\int_{\mathbb{T}} h_\epsilon(y)\Big(\sum_{\substack{0<\gamma\le T\\ \{\alpha\gamma\}\in\mathbb{I}+y}} 1 - N(T)|\mathbb{I}|\Big)\,dy = T\int_0^{\varepsilon} h_\epsilon(y)\int_{\mathbb{I}+y} g_\alpha(x)dx\,dy + o(T). \tag{2.1}$$

By (1.3), g_α is bounded and it follows that

$$\Big|\int_{\mathbb{I}+y} g_\alpha(x)dx - \int_{\mathbb{I}} g_\alpha(x)dx\Big| \ll \epsilon$$

for $0 \le y \le \varepsilon$. Therefore the right side of (2.1) equals

$$T \int_{\mathbb{I}} g_\alpha(x)dx + o(T) + O(\epsilon T).$$

It follows that for some choice of $y \in (0, \epsilon)$ one must have

$$\Big| \sum_{\substack{0<\gamma\le T \\ \{\alpha\gamma\}\in\mathbb{I}+y}} 1 - N(T)|\mathbb{I}| \Big| \ge T \Big| \int_{\mathbb{I}} g_\alpha(x)dx \Big| + o(T) + O(\epsilon T).$$

Letting $\epsilon \to 0$, we obtain our lower bound for the discrepancy.

Proof (Proof of Theorem 1 (ii) and Theorem 2). Let

$$h(u) = \begin{cases} 1 & \{u\} \in \mathbb{I} \\ 0 & \text{else} \end{cases}$$

and let J be a positive integer. There are trigonometric polynomials h^+ and h^-, depending on J and $\mathbb{I}$, satisfying

$$\begin{aligned} h^-(u) \le h(u) \le h^+(u) &\qquad (u \in \mathbb{R}), \\ h^\pm(u) = \sum_{|j|\le J} c_j^\pm e^{2\pi i j u}&, \\ c_0^\pm = |\mathbb{I}| \pm \frac{1}{J+1}, &\qquad |c_j^\pm| \le \frac{1}{|j|} \quad (j \ge 1). \end{aligned}$$

For proofs, see Chapter 1 of [15], for example. These trigonometric polynomials are optimal in the sense that with J fixed, $|c_0^\pm - |\mathbb{I}||$ cannot be made smaller. We have

$$\sum_{0<\gamma\le T} h^-(\alpha\gamma) \le \sum_{\substack{0<\gamma\le T \\ \{\alpha\gamma\}\in\mathbb{I}}} 1 \le \sum_{0<\gamma\le T} h^+(\alpha\gamma).$$

For integers j, let $x_j = e^{2\pi j\alpha}$ and for positive j put

$$V_j = \frac{-\Lambda(n_{x_j})}{2\pi x_j^{1/2}} \frac{e^{iT\log(x_j/n_{x_j})} - 1}{i\log(x_j/n_{x_j})}.$$

Also define $V_{-j} = \overline{V_j}$. By (1.6), for nonzero j we have

$$\sum_{0<\gamma\le T} x_j^{i\gamma} = V_j + O\left(x_{|j|}^{1/2}\log^2(x_{|j|}T)\right).$$

This will be used for

$$1 \le |j| \le J_0 := \left\lfloor \frac{2\log T - 5\log\log T}{2\pi\alpha} \right\rfloor.$$

Suppose that $J \ge J_0$. We obtain (implied constants depend on α)

$$\begin{aligned}\sum_{0<\gamma\le T} h^{\pm}(\alpha\gamma) &= c_0^{\pm} N(T) + \sum_{1\le|j|\le J} c_j^{\pm} \sum_{0<\gamma\le T} x_j^{i\gamma} \\ &= c_0^{\pm} N(T) + 2\Re \sum_{1\le j\le J_0} c_j^{\pm} \Big[V_j + O(x_j^{1/2}\log^2 T) \Big] \\ &\quad + \sum_{J_0<|j|\le J} O\left(\frac{1}{|j|}\right) \Big| \sum_{0<\gamma\le T} x_j^{i\gamma} \Big| \\ &= |\mathbb{I}| N(T) + \sum_{j\ne 0} c_j^{\pm} V_j \pm \frac{N(T)}{J+1} + o(T) + \sum_{J_0<|j|\le J} O(|j|^{-1}) \Big| \sum_{0<\gamma\le T} x_j^{i\gamma} \Big|.\end{aligned}$$

If $\alpha = \frac{a}{q}\frac{\log p}{2\pi}$ for a prime p and coprime positive a, q, then $x_j = p^{aj/q}$ and consequently

$$V_{kq} = -\frac{T\log p}{2\pi p^{ak/2}}$$

for nonzero integers k. Thus,

$$\sum_{\substack{j\ne 0 \\ q|j}} c_j^{\pm} V_j = T \int_{\mathbb{T}} h^{\pm} g_\alpha.$$

If $q \nmid j$, then x_j is not an integer. Hence

$$\sum_{\substack{j\ne 0 \\ q\nmid j}} c_j^{\pm} V_j \ll T \sum_{\substack{1\le|j|\le J \\ q\nmid j}} \frac{1}{e^{\pi j\alpha}} \left| \frac{e^{iT\log(x_j/n_{x_j})} - 1}{iT\log(x_j/n_{x_j})} \right|.$$

The sum on the right converges uniformly in T, and each summand is $o(1)$ as $T\to\infty$, hence the left side is $o(T)$. We conclude

$$\sum_{j\ne 0} c_j^{\pm} V_j = T \int_{\mathbb{T}} h^{\pm} g_\alpha + o(T). \tag{2.2}$$

When α is not of the form $\frac{a}{q}\frac{\log p}{2\pi}$, x_j is never an integer (for nonzero j), and a similar argument yields (2.2). Since $h - h^{\pm}$ has constant sign,

$$\Big| \int_{\mathbb{T}} (h - h^{\pm}) g_\alpha \Big| \le \max_{x\in\mathbb{T}} |g_\alpha(x)| \int_{\mathbb{T}} |h - h^{\pm}| = \frac{\max_{x\in\mathbb{T}} |g_\alpha(x)|}{J+1} \ll \frac{1}{\log T}.$$

Therefore,

$$\sum_{0<\gamma\le T} h^{\pm}(\alpha\gamma) = |\mathbb{I}|N(T) + T\int_{\mathbb{T}} hg_\alpha + o(T) \pm \frac{N(T)}{J+1} + \sum_{J_0<|j|\le J} O\left(\frac{1}{|j|}\right)\Big|\sum_{0<\gamma\le T} x_j^{i\gamma}\Big|.$$

For Theorem 1 (ii), we take $J = J_0$. For Theorem 2, take $J = \lfloor \lambda \log T \rfloor$ with λ fixed, and then let $\lambda \to \infty$.

Proof (Proof of Theorem 3). We first construct a function F which is a good approximation of the characteristic function of the interval $[0,1]$ and whose Fourier transform is supported on $[-K,K]$, where K is a parameter to be specified later. Consider the entire function

$$H(z) = \left(\frac{\sin \pi z}{\pi}\right)^2 \left(\sum_{n=1}^{\infty} \frac{1}{(z-n)^2} - \sum_{n=1}^{\infty} \frac{1}{(z+n)^2} + \frac{2}{z}\right)$$

for complex z, and set

$$F(z) = \frac{H(Kz) + H(K - Kz)}{2}.$$

The function $H(z)$ is related to the so-called Beurling-Selberg functions, and basic facts about H can be found in [23]. In particular, for real x, (i) $H(x)$ is an odd function; (ii) the Fourier transform $\widehat{H}$ is supported on $[-1,1]$; (iii) $H(x) = \mathrm{sgn}(x) + O(\frac{1}{1+|x|^3})$, where $\mathrm{sgn}(x) = 1$ if $x > 0$, $\mathrm{sgn}(x) = -1$ if $x < 0$ and $\mathrm{sgn}(0) = 0$; (iv) $H'(x) = O(\frac{1}{1+|x|^3})$. Item (iii) follows from (2.26) of [23] and the Euler-Maclaurin summation formula, and (iv) follows from Theorem 6 of [23]. Let I be the indicator function of the interval $[0,1]$. It follows that the Fourier transform $\widehat{F}$ of F is supported on $[-K,K]$ and

$$|F(x) - I(x)| \ll \frac{1}{1+K^3|x|^3} + \frac{1}{1+K^3|1-x|^3}. \tag{2.3}$$

Since

$$\widehat{I}(t) = \frac{1 - e^{-2\pi i t}}{2\pi i t},$$

it follows readily that $\widehat{F}(t) \ll 1$, uniformly in K, and

$$\widehat{F}'(t) = \frac{1-(1+2\pi it)e^{-2\pi it}}{-2\pi it^2} + O\bigg(\int_{-\infty}^{\infty} \frac{|x|}{1+K^3|x|^3} + \frac{|x|}{1+K^3|1-x|^3}\,dx\bigg)$$
$$= O\left(\frac{1}{1+|t|} + \frac{1}{K}\right).$$

Next, let $T \ge 2$ and $T \le x \le T^A$. Write

$$\sum_{0<\gamma\le T} x^{i\gamma} = \sum_{|\gamma|\le x} x^{i\gamma} F(\gamma/T) + \sum_{|\gamma|\le x} x^{i\gamma}\big[I(\gamma/T) - F(\gamma/T)\big].$$

By (1.2) and (2.3), the second sum on the right is

$$\le 2N\left(\frac{T}{K}\right) + N\left(T+\frac{T}{K}\right) - N\left(T-\frac{T}{K}\right)$$
$$+ \frac{T^3}{K^3}\bigg(\sum_{|\gamma|>T/K} \frac{1}{|\gamma|^3} + \sum_{|\gamma-T|\ge T/K} \frac{1}{|\gamma-T|^3}\bigg)$$
$$\ll \frac{T\log T}{K}.$$

Also,

$$\sum_{|\gamma|\le x} x^{i\gamma}F(\gamma/T) \sum_{|\gamma|\le x} x^{i\gamma} \int_{-K}^{K} e^{2\pi iv\gamma/T}\widehat{F}(v)\,dv$$
$$= x^{-1/2}\int_{-K}^{K} e^{-\pi v/T}\widehat{F}(v) \sum_{|\gamma|\le x} \left(xe^{2\pi v/T}\right)^{\rho} dv$$
$$= -\frac{T}{2\pi x^{1/2}} \int_{-K}^{K} e^{-\pi v/T}\left(\widehat{F}'(v) - \frac{\pi}{T}\widehat{F}(v)\right) \sum_{|\gamma|\le x} \frac{\left(xe^{2\pi v/T}\right)^{\rho}}{\rho}\,dv,$$

where the last line follows from the previous line using integration by parts. The final sum on γ is evaluated using the explicit formula (see e.g. [2], §17)

$$G(x) := \psi(x) - x = -\sum_{|\gamma|\le M} \frac{x^{\rho}}{\rho} + O\left(\log x + \frac{x\log^2(Mx)}{M}\right), \qquad (2.4)$$

valid for $x \ge 2$, $M \ge 2$. Since

$$\int_{-K}^{K} e^{-\pi v/T}\left(\widehat{F}'(v) - \frac{\pi}{T}\widehat{F}(v)\right) dv = 0,$$

we obtain

$$\sum_{|\gamma|\le x} x^{i\gamma}F(\gamma/T) = \frac{-T}{2\pi\sqrt{x}}\int_{-K}^{K}\widehat{F}'(v)\left(G(xe^{2\pi v/T})-G(x)\right)\,dv + O\left(K\left(1+Tx^{-1/2}\right)\log^2 x\right).$$

Altogether, this gives

$$\sum_{|\gamma|\le T} x^{i\gamma} \ll \frac{T\log K}{\sqrt{x}}\max_{xe^{-2\pi K/T}\le y\le xe^{2\pi K/T}}|G(y)-G(x)| + \frac{T\log T}{K} + K\left(1+Tx^{-1/2}\right)\log^2 x.$$

Take $K=\log^2 T$ and assume Conjecture 4. The first part of Theorem 3 follows.

The second part is straightforward, starting with the explicit formula (2.4) in the form

$$\psi(x+y)-\psi(x)-y = -\sum_{|\gamma|\le x}\frac{(x+y)^\rho - x^\rho}{\rho} + O(\log^2 x).$$

Fix $\varepsilon>0$ and apply Conjecture 3 with $A=2/\varepsilon$. By partial summation,

$$\begin{aligned}\Big|\sum_{x^{\varepsilon/2}<|\gamma|\le x}\frac{x^\rho}{\rho}\Big| &= 2\Big|\Re\sum_{x^{\varepsilon/2}<\gamma\le x}\frac{x^\rho}{\rho}\Big| \\ &\le 2x^{1/2}\left[\frac{1}{\frac12+ix}\sum_{0<\gamma\le x}x^{i\gamma} + i\int_{x^{\varepsilon/2}}^{x}\frac{1}{(\frac12+it)^2}\sum_{0<\gamma\le t}x^{i\gamma}\,dt\right] \\ &= o(x^{1/2}\log x).\end{aligned}$$

The smaller zeros are handled in a trivial way. We have, for $y\le x$,

$$(x+y)^\rho - x^\rho = x^\rho\left(\rho\frac{y}{x}+O\left(\frac{|\rho|^2y^2}{x^2}\right)\right),$$

whence

$$\sum_{|\gamma|\le x^{\varepsilon/2}}\frac{(x+y)^\rho-x^\rho}{\rho} \ll N(x^{\varepsilon/2})x^{1/2}\left(\frac{y}{x}+x^{\varepsilon/2}\frac{y^2}{x^2}\right) \ll x^{\frac12-\frac{\varepsilon}{2}}\log x.$$

Therefore, $\psi(x+y)-\psi(x)-y=o(x^{1/2}\log x)$, as claimed.

Proof (Proof of Theorem 4). It will be convenient to work with the normalized sum

$$\mathcal{D}(x,T)=\frac{\mathcal{F}(x,T)}{N(T)}.$$

Lemma 1. *Suppose $T\ge 10$ and $1\le\beta\le T$. Then*

$$\begin{aligned}\sum_{0<\gamma\le T}x^{i\gamma}&\ll T\left(\frac{\log T}{\beta}\right)^{\frac12}\Bigg(1+\max_{\frac{T}{\beta\log T}\le t\le T}|\mathcal{D}(x,t)|\\&\qquad+\beta^3\left|\int_{-\infty}^{\infty}(\mathcal{D}(xe^u,t)-\mathcal{D}(x,t))e^{-2\beta|u|}\,du\right|\Bigg)^{\frac12}\\&\ll\frac{T(\log T)^{\frac12}}{\beta^{1/2}}\left(1+\max_{\frac{T}{\beta\log T}\le t\le T}|\mathcal{D}(x,t)|\right)^{1/2}+T(\beta\log T)^{1/2}\\&\quad\times\left(\max_{\frac{T}{\beta\log T}\le t\le T}\ \max_{0\le u\le\frac{1}{\beta}\log(\beta\log T)}|\mathcal{D}(xe^u,t)+\mathcal{D}(xe^{-u},t)-2\mathcal{D}(x,t)|\right)^{\frac12}.\end{aligned}$$

Proof. We follow [6] by estimating $\sum_{0<\gamma\le T}x^{i\gamma}$ in terms of

$$G_\beta(x,T)=\sum_{0<\gamma,\gamma'\le T}\frac{4\beta^2x^{i(\gamma-\gamma')}}{4\beta^2+(\gamma-\gamma')^2}.$$

In particular, $G_1(x,T)=\mathcal{F}(x,T)$, and by (1.2), we have $G_\beta(x,T)\ll(1+\beta)T\log^2T$. By Lemma 1 of [6], uniformly for $1\le\beta\le T$ and $1\le V\le T$, we have

$$\begin{aligned}\sum_{0<\gamma\le T}x^{i\gamma}&\ll\Big(T\beta^{-1}\max_{t\le T}G_\beta(x,t)\Big)^{1/2}\\&\ll\frac{T\log T}{V^{1/2}}+\Big(T\beta^{-1}\max_{T/V\le t\le T}G_\beta(x,t)\Big)^{1/2}.\end{aligned}\tag{2.5}$$

Using Lemma 2 of [6], we have

$$\begin{aligned}G_\beta(x,t)&=\beta^2\mathcal{F}(x,t)+\beta(1-\beta^2)\int_{-\infty}^{\infty}\mathcal{F}(xe^u,t)e^{-2\beta|u|}\,du\\&=\mathcal{F}(x,t)+\beta(1-\beta^2)\int_{-\infty}^{\infty}(\mathcal{F}(xe^u,t)-\mathcal{F}(x,t))e^{-2\beta|u|}\,du,\end{aligned}$$

from which the first inequality in the lemma follows upon taking $V=\beta\log T$. For the second inequality, combine the terms in the integral with $u=v$ and $u=-v$ for $0\le v\le\frac{\log(\beta\log T)}{\beta}$, and use the trivial bound $\mathcal{D}(z,t)\ll\log t$ when $|u|\ge\frac{\log(\beta\log T)}{\beta}$ ($z=x$ and $z=xe^u$).

In order to finish the proof of Theorem 4, suppose that $\log T \le \beta \le \log^2 T$. From Conjecture 5 it follows that the terms $\mathcal{D}(xe^u, t)$, $\mathcal{D}(xe^{-u}, t)$, and $\mathcal{D}(x, t)$, in the ranges from the statement of the above lemma, are all of the form $1 + o\left((\log T)^{-2}\right)$. Therefore,

$$\sum_{0<\gamma\le T} x^{i\gamma} = O\left(T\frac{(\log T)^{1/2}}{\beta^{1/2}}\right) + o\left(T\frac{\beta^{1/2}}{(\log T)^{1/2}}\right).$$

Thus, taking β slightly larger than $\log T$ produces the desired result.

3. General *L*-functions

Consider a Dirichlet series $F(s) = \sum_{n=1}^{\infty} a_F(n)n^{-s}$ satisfying the following axioms:
(i) there exists an integer $m \ge 0$ such that $(s-1)^m F(s)$ is an entire function of finite order;
(ii) F satisfies a functional equation of the type:

$$\Phi(s) = w\overline{\Phi}(1-s),$$

where

$$\Phi(s) = Q^s \prod_{j=1}^{r} \Gamma(\lambda_j s + \mu_j) F(s)$$

with $Q > 0$, $\lambda_j > 0$, $\Re(\mu_j) \ge 0$ and $|w| = 1$. (Here, $\overline{f}(s) = \overline{f(\overline{s})}$);
(iii) $F(s)$ has an Euler product, which we write as

$$-\frac{F'}{F}(s) = \sum_{n=1}^{\infty} \Lambda_F(n) n^{-s},$$

where $\Lambda_F(n)$ is supported on powers of primes.

We also need some growth conditions on the coefficients $a_F(n)$ and $\Lambda_F(n)$. Although stronger than we require, for convenience we impose the conditions (iv) $\Lambda_F(n) \ll n^{\theta_F}$ for some $\theta_F < \frac{1}{2}$ and (v) for every $\varepsilon > 0$, $a_F(n) \ll_\varepsilon n^\varepsilon$. Together, conditions (i)–(v) define the *Selberg class* $\mathcal{S}$ of Dirichlet series. For a survey of results and conjectures concerning the Selberg class, the reader may consult Kaczorowski and Perelli's paper [10]. In particular, $\mathcal{S}$ includes the Riemann zeta function, Dirichlet L-functions, and L-functions attached to number fields and elliptic curves. The Selberg class is conjectured to equal the class of all automorphic L-functions, suitably normalized so that their nontrivial zeros have real parts between 0 and 1.

The functional equation is not uniquely determined in light of the duplication formula for Γ-function, however the real sum

$$d_F = 2\sum_{j=1}^{r} \lambda_j$$

is well-defined and is known as the degree of F. Analogous to (1.2), we have (cf. [22], (1.6))

$$\begin{aligned} N_F(T) &= |\{\rho = \beta + i\gamma : F(\rho) = 0, 0 < \beta < 1, 0 < \gamma \le T\}| \\ &= \frac{d_F}{2\pi} T \log T + c_1 T + O(\log T) \end{aligned} \tag{3.1}$$

for some constant $c_1 = c_1(F)$. A function $F \in \mathcal{S}$ is said to be *primitive* if it cannot be written as a product of two or more elements of $\mathcal{S}$. Clearly it suffices to prove our results in the case of F being primitive, which we henceforth assume. It is expected that all zeros of F with real part between 0 and 1 have real part $\frac{1}{2}$, a hypothesis we abbreviate as RH_F. Although we shall assume RH_F for many of the results in this section, sometimes a weaker hypothesis suffices, that most zeros of F are close to the critical line.

Hypothesis Z_F. There exist constants $A > 0, B > 0$ (depending on F) such that

$$\begin{aligned} N_F(\sigma, T) &= \left| \left\{ \beta + i\gamma : \frac{1}{2} \le \beta \le \sigma, 0 < \gamma \le T \right\} \right| \\ &\ll T^{1 - A(\sigma - 1/2)} \log^B T, \end{aligned}$$

uniformly for $\sigma \ge 1/2$ and $T \ge 2$.

Hypothesis Z_F is known, with $B = 1$, for the Riemann zeta function and Dirichlet L-functions (Selberg [20], [21]), and certain degree 2 L-functions attached to cusp forms (Luo [13]).

The next tool we require is an analog of (1.6). It is very similar to Proposition 1 of [19], and with small modifications to that proof we obtain the following result, which is nontrivial provided $x^{1/2+\theta_F} + x^{1/2+\varepsilon} \ll T$.

Lemma 2. *Let $F \in \mathcal{S}$, $x > 1$, $T \ge 2$, and let n_x be a nearest integer to x. Then, for any $\varepsilon > 0$,*

$$\begin{aligned} \sum_{0 < \gamma \le T} x^\rho = &- \frac{\Lambda_F(n_x)}{2\pi} \frac{e^{iT\log(x/n_x)} - 1}{i \log(x/n_x)} \\ &+ O_\varepsilon \left(x^{1+\theta_F} \log(2x) + x^{1+\varepsilon} \log T + \frac{\log T}{\log x} \right). \end{aligned}$$

Using Lemma 2 in place of Lemma 1 of [3], Hypothesis Z_F in place of Lemma 2 of [3], and following the proof of Theorem 1 of [3], we obtain a generalization of (1.4).

Theorem 5. *Let $F \in \mathcal{S}$. If $\alpha = \frac{a \log p}{2\pi q}$ for some prime number p and positive integers a, q with $(a,q) = 1$, define*

$$g_{F,\alpha}(t) = -\frac{1}{\pi} \Re \sum_{k=1}^{\infty} \frac{\Lambda_F(p^{ak})}{p^{ak/2}} e^{-2\pi i q k t},$$

For other α, define $g_{F,\alpha}(t) = 0$ for all t. If Hypothesis Z_F holds, then

$$\sum_{0<\gamma\le T} f(\alpha\gamma) = N_F(T) \int_{\mathbb{T}} f(x)\,dx + T \int_{\mathbb{T}} f(x) g_{F,\alpha}(x)\,dx + o(T) \qquad (3.2)$$

for all $f \in C^2(\mathbb{T})$. Assuming RH_F, (3.2) holds for all absolutely continuous f.

Since Hypothesis Z_F holds for Dirichlet L-functions $L(s,\chi)$, we obtain the following.

Corollary 1. *Unconditionally, for Dirichlet L-functions F, (3.2) holds for all $f \in C^2(\mathbb{T})$.*

When $F(s) = L(s,\chi)$ and $\alpha = \frac{a \log p}{2\pi q}$ with p prime, $(a,q) = 1$, we have


$$g_{F,\alpha}(t) = -\frac{\log p}{\pi} \Re \left(\frac{e^{2\pi i(qt + a\xi)}}{p^{a/2} - e^{2\pi i(qt+a\xi)}} \right),$$

where $\chi(p) = e^{2\pi i \xi}$. It follows that there is a shortage of zeros of $L(s,\chi)$ with $\{\alpha\gamma\}$ near $\frac{k - a\xi}{q}$, $k = 0, \cdots, q-1$. We illustrate this phenomenon with three histograms of $M_F(y;T)$, F a Dirichlet L-function associated with a character of conductor 5 and $T = 500,000$. For both characters, $N_F(T) = 946488$. The list of zeros was taken from Michael Rubinstein's data files on his Web page. In Figure 1 we plot for each subinterval $I = [y, y + \frac{1}{500})$ the value of $500(M_F(y + \frac{1}{500}) - M_F(y))$ and also the graph of $g_{F,\alpha}(y)$. Here


$$M_F(y) = \frac{T}{N_F(T)} \left| \sum_{\substack{0<\gamma\le T \\ \{\alpha\gamma\}<y}} 1 - y N_F(T) \right|$$

The characters are identified by their value at 2.


We conjecture that (3.2) holds when f is the indicator function of an interval, and are thus led to the following generalizations of Conjectures 1 and 2. Here $D_{F,\alpha}$ is the natural generalization of the discrepancy function D_α.



$\alpha = \frac{\log 2}{2\pi}, \chi(2) = -i$



$\alpha = \frac{\log 3}{2\pi}, \chi(2) = -i$



$\alpha = \frac{\log 3}{2\pi}, \chi(2) = -1$

Fig. 1. $500(M_F(y + \frac{1}{500}) - M_F(y))$ vs. $g_{F,\alpha}(y)$ for $T = 500000$.

Conjecture 6. Let $\mathbb{I}$ be an interval of $\mathbb{T}$. Then

$$\sum_{\substack{0<\gamma\le T\\ \{\alpha\gamma\}\in\mathbb{I}}} 1 = |\mathbb{I}|N_F(T) + T\int_{\mathbb{I}} g_{F,\alpha}(x)dx + o(T).$$

Conjecture 7. We have

$$D_{F,\alpha}(T) = \frac{T}{N_F(T)}\sup_{\mathbb{I}}\Big|\int_{\mathbb{I}} g_{F,\alpha}(x)\,dx\Big| + o\Big(\frac{1}{\log T}\Big).$$

Combining Theorem 5 and the proof of Theorem 1, we obtain the following. The only difference in the proof is that here we take

$$J_0 = \left\lfloor \frac{\frac{\log T}{1/2+\theta_F} - 5\log\log T}{2\pi\alpha} \right\rfloor.$$

Theorem 6. *(i) Assuming Hypothesis Z_F, we have*

$$D_{F,\alpha}(T) \ge \frac{T}{N_F(T)}\sup_{\mathbb{I}}\Big|\int_{\mathbb{I}} g_{F,\alpha}(x)\,dx\Big| + o\Big(\frac{1}{\log T}\Big).$$

(ii) Assuming RH_F, for any interval $\mathbb{I}$ of $\mathbb{T}$ we have

$$\Big|\sum_{\substack{0<\gamma\le T\\ \{\alpha\gamma\}\in\mathbb{I}}} 1 - |\mathbb{I}|N_F(T) - T\int_{\mathbb{I}} g_{F,\alpha}(x)dx\Big| \le \alpha(1/2+\theta_F)T + o(T).$$

We can prove a direct analog of Theorem 2, by requiring a slightly larger range of T in the analog of Conjecture 3, since θ_F may be large.

Conjecture 8. Let $A > 1$ be a fixed real number. Uniformly for

$$\frac{T^{1/(1/2+\theta_F)}}{\log^5 T} \le x \le T^A,$$

we conjecture that

$$\sum_{0<\gamma\le T} x^{i\gamma} = o(T). \tag{3.3}$$

Theorem 7. *Assume RH_F. Then Conjecture 8 implies Conjectures 6 and 7.*

The analog of Theorem 3 holds for $F \in \mathcal{S}$, by following the proof given in the preceding section. Here we need an explicit formula similar to (2.4). By standard contour integration methods, one obtains

$$G_F(x) := \sum_{n \le x} \Lambda_F(n) - d_F x = -\sum_{|\rho| \le Q} \frac{x^\rho}{\rho} + O(x^{\theta_F} \log x)$$

provided $Q \ge x \log x$. Since $\theta_F < \frac{1}{2}$, the error term is acceptable.

Conjecture 9. For every $\varepsilon > 0$, if x is large and $y \le x^{1-\epsilon}$, then

$$G_F(x+y) - G_F(x) = o(x^{\frac{1}{2}} / \log\log x).$$

Theorem 8. *Assume RH_F. Conjecture 9 implies Conjecture 8, and hence Conjectures 6 and 7. Conversely, if RH_F and Conjecture 8 holds, then for all fixed $\varepsilon > 0$, large x and $y \le x^{1-\varepsilon}$,*

$$G_F(x+y) - G_F(x) = o(x^{\frac{1}{2}} \log x).$$

In order to address an analog of Theorem 4, we first quote a Pair Correlation Conjecture for F, due to Murty and Perelli [18].

Conjecture 10. Define

$$\mathcal{F}_F(x,T) = \sum_{0<\gamma,\gamma' \le T} \frac{4x^{i(\gamma-\gamma')}}{4+(\gamma-\gamma')^2}$$

and $\mathcal{D}_F(x,T) = \mathcal{F}_F(x,T)/N_F(T)$. We have $\mathcal{D}_F(T^{\theta d_F},T) \sim \theta$ for $0 < \theta \le 1$ and $\mathcal{D}(T^{\theta d_F},T) \sim 1$ for $\theta \ge 1$.

Notice that, as a function of x, $\mathcal{F}_F(x,T)$ is conjectured to undergo a change of behavior in the vicinity of $x = T^{d_F}$. In order to deduce Conjecture 8, we can postulate a stronger version of Conjecture 10, with error terms of relative order $o(1/\log^2 T)$. We succeed, as in the proof of Theorem 4, when $d_F = 1$. When $d_F \ge 2$, however, this transition zone lies outside the range in which Lemma 2 is useful (Kaczorowski and Perelli recently proved that $1 < d_F < 2$ is impossible [11]; it is conjectured that d_F is always an integer). We can use an analog of Lemma 2, which follows by the same method (replace $\mathcal{D}(x,T)$ with $\mathcal{D}_F(x,T)$). However, in order to prove the right side is small, we require that $\mathcal{D}_F(x,T)$ has small *variation*, even through the transition zone $x \approx T^{d_F}$. Tsz Ho Chan [1] studied the behavior of $\mathcal{D}(x,T)$ (for $\zeta(s)$) in the vicinity of $x = T$ assuming RH plus a quantitative version of the twin prime conjecture with strong error term. His analysis leads to a pair correlation conjecture with $\mathcal{D}(x,T)$ smoothly varying through the transition zone. We conjecture that the same holds for other $F \in \mathcal{S}$.

Conjecture 11. Suppose $F \in \mathcal{S}$, $\mathcal{D}_F(x,T) \ll 1$ uniformly in x and T, and for any $A > 0$ there is a $c > 0$ so that

$$|\mathcal{D}_F(x+\delta x,T) + \mathcal{D}_F(x-\delta x,T) - 2\mathcal{D}_F(x,T)| = o(T/\log T)$$

uniformly for $T \le x \le T^A$ and $0 \le \delta \le (\log T)^{c-1}$.

Following the proof of Theorem 4 (take $\beta = \log T \log\log T$, for example), we arrive at the following.

Theorem 9. *Assume RH_F. Then Conjecture 11 implies Conjecture 8, and therefore also Conjectures 6 and 7.*

Acknowledgement. The authors thank the referee for carefully reading the paper and for pointing out several misprints and minor errors.

References

1. T. H. Chan, *More precise pair correlation conjecture on the zeros of the Riemann zeta function*, Acta Arith. **114** (2004), no. 3, 199–214.
2. H. Davenport, *Multiplicative Number Theory*, 3rd ed., Springer-Verlag, 2000.
3. K. Ford and A. Zaharescu, *On the distribution of imaginary parts of zeros of the Riemann zeta function*, J. Reine Angew. Math. **579** (2005), 145–158.
4. A. Fujii, *On the zeros of Dirichlet L-functions, III*, Trans. Amer. Math. Soc. **219** (1976), 347–349.
5. P. X. Gallagher and J. H. Mueller, *Primes and zeros in short intervals*, J. Reine Angew. Math. **303/304** (1978), 205–220.
6. D. A. Goldston and D. R. Heath-Brown, *A note on the differences between consecutive primes*, Math. Ann. **266** (1984), 317–320.
7. S. M. Gonek, *An explicit formula of Landau and its applications to the theory of the zeta-function,* A tribute to Emil Grosswald: number theory and related analysis, 395–413, Contemp. Math., 143, Amer. Math. Soc., Providence, RI, 1993.
8. D. R. Heath-Brown, *Gaps between primes, and the pair correlation of zeros of the zeta-function*, Acta Arith. **41** (1982), 85–99.
9. E. Hlawka, *Über die Gleichverteilung gewisser Folgen, welche mit den Nullstellen der Zetafunktionen zusammenhängen,* Sitzungsber. Österr. Akad. Wiss., Math.–Naturnw. Kl. Abt. II **184** (1975), 459–471.
10. J. Kaczorowski and A. Perelli, *The Selberg class: a survey*, Number Theory in Progress, vol. II, de Gruyter, Berlin (1999), 953–992.
11. J. Kaczorowski and A. Perelli, *Nonexistence of L-functions of degree $1 < d < 2$*, preprint.
12. E. Landau. *Über die Nullstellen der ζ-Funktion*, Math. Ann. **71** (1911), 548–568.
13. W. Luo, *Zeros of Hecke L-functions associated with cusp forms.*, Acta Arith. **71** (1995), no. 2, 139–158.
14. H. L. Montgomery, *The pair correlation of zeros of the zeta function,* Proc. Sym. Pure Math. **24** (1973), 181–193.

15. H. L. Montgomery, *Ten lectures on the interface between analytic number theory and harmonic analysis*. CBMS Regional Conference Series in Mathematics, 84. American Mathematical Society, Providence, RI, 1994. xiv+220 pp.
16. H. L. Montgomery and K. Soundararajan, *Primes in short intervals*, Commun. Math. Phys. **252** (2004), 589–617.
17. J. H. Mueller, *On the difference between consecutive primes*, Recent progress in analytic number theory, I, pp. 269–273. London, New York: Academic Press 1981.
18. M. R. Murty, A. Perelli, *The Pair Correlation of Zeros of Functions in the Selberg Class*, Int. Math. Res. Not. (1999) No. **10**, 531–545.
19. M. R. Murty, A. Zaharescu, *Explicit formulas for the pair correlation of zeros of functions in the Selberg class*, Forum Math. **14** (2002), no. 1, 65–83.
20. A. Selberg, *Contributions to the theory of the Riemann zeta-function*, Arch. Math. Naturvid. **48** (1946), 89–155; Collected papers, vol. I, 214–280, Springer, Berlin 1989.
21. A. Selberg, *Contributions to the theory of Dirichlet's L-functions*, Skr. Norske Vid. Akad. Oslo. I. **1946**, (1946), no. 3, 62 pp.; Collected papers, vol. I, 281–340, Springer, Berlin 1989.
22. A. Selberg, *Old and new conjectures and results about a class of Dirichlet series*, Proceedings of the Amalfi Conference on Analytic Number Theory (Maiori, 1989), Univ. Salerno, (1992), 367–385; Collected papers, vol. II, 47–63, Springer, Berlin 1989.
23. J. Vaaler, *Some extremal functions in Fourier analysis*, Bull. Amer. Math. Soc. (N.S.) **12** (1985), no. 2, 183–216.