arXiv:2007.03474v5 [math.AT] 4 Feb 2021

7-DIMENSIONAL SIMPLY-CONNECTED SPIN MANIFOLDS WHOSE INTEGRAL COHOMOLOGY RINGS ARE ISOMORPHIC TO THAT OF $\mathbb{C}P^2 \times S^3$ ADMIT ROUND FOLD MAPS

NAOKI KITAZAWA

Abstract. 7-dimensional closed and simply-connected manifolds are important objects in the theory of classical algebraic topology and differential topology. The class has been attractive since the discoveries of 7-dimensional exotic homotopy spheres by Milnor and so on. Still recently, new understandings via algebraic topological tools such as characteristic classes and bordism relations have been discovered by Kreck and so on.

The author has been interested in understanding the class in geometric and constructive ways. The author constructed explicit *fold* maps, which are higher dimensional versions of Morse functions, on the manifolds. The studies have been motivated by studies of *special generic* maps, which are higher dimensional versions of Morse functions on homotopy spheres with exactly two singular points, characterizing them topologically except 4-dimensional cases. The class contains canonical projections of unit spheres for example. This class has been found to be interesting, restricting the topologies and the differentiable structures of the manifolds strictly: Saeki, Sakuma, Wrazidlo, and so on, found explicit phenomena. The present paper concerns fold maps on 7-dimensional closed, simply-connected and spin manifolds whose integral cohomology rings are isomorphic to that of $\mathbb{C}P^2 \times S^3$.

1. Introduction, terminologies and notation.

7-dimensional closed and simply-connected manifolds are important objects in the theory of classical algebraic topology and differential topology (of higher dimensional closed and simply-connected manifolds). The class has been attractive since the discoveries of 7-dimensional exotic homotopy spheres by Milnor and so on. Still recently, new understandings via algebraic topological tools such as characteristic classes and bordism relations have been performed by Kreck [9] ,Wang [18] and so on.

The author has been interested in understanding the class in geometric and constructive ways and obtained explicit *fold* maps, which are higher dimensional versions of Morse functions, on the manifolds. The studies have been motivated by studies of *special generic* maps, which are higher dimensional versions of Morse functions on homotopy spheres with exactly two singular points, characterizing them topologically except 4-dimensional cases and the class contains canonical projections of unit spheres for example. This class has been found to be interesting, restricting the topologies and the differentiable structures of homotopy spheres and manifolds admitting them strictly. For example 7-dimensional exotic homotopy

2020 *Mathematics Subject Classification.* Primary 57R45. Secondary 57R19.

Key words and phrases. Singularities of differentiable maps; fold maps. Cohomology rings. Higher dimensional closed and simply-connected manifolds.

spheres, which are not diffeomorphic to the standard sphere S^7, admit no special geneirc map into $\mathbb{R}^n$ for $n = 4, 5, 6$ and in considerable cases admit no special generic maps for $n = 3$ ([14], [15], [16], [17], [19] and so on). It is also an important fact that construction of explicit fold maps on explicit manifolds is difficult even for fundamental manifolds and in general. The present paper concerns fold maps on 7-dimensional simply-connected spin manifolds whose integral cohomology rings are isomorphic to that of $\mathbb{C}P^2 \times S^3$. A recent study on classifications of these manifolds by [18] motivates the author to do this study.

1.1. **Terminologies and notation on smooth maps and so on.** Throughout the present paper, manifolds and maps between manifolds are smooth (of class C^∞). Diffeomorphisms on manifolds are always smooth and the *diffeomorphism group* of a manifold is defined as the group of all smooth diffeomorphisms on the manifold. For bundles whose fibers are manifolds, the structure groups are subgroups of the diffeomorphism groups or the bundles are assumed to be *smooth* unless otherwise stated. A *linear* bundle means a smooth bundle whose fiber is a unit sphere or a unit disc in a Euclidean space with structure group acting linearly in a canonical way on the fiber.

A *singular* point of a smooth map is a point in the domain at which the dimension of the image of the differential is smaller than both the dimensions of the domain and the target. We call the set of all singular points the *singular set* of the map. We call the image of the singular set the *singular value set* of the map. The *regular value set* of the map is the complementary set of the singular value set. A *singular* (*regular*) *value* is a point in the singular (resp. regular) value set. For a smooth map c, $S(c)$ denotes the singular set of c.

$||x||$ denotes the distance between $x \in \mathbb{R}^k$ and the origin $0 \in \mathbb{R}^k$ where the Euclidean space is endowed with the standard metric.

1.2. **The definition and fundamental properties of a fold map and explicit fold maps.** Hereafter, let $m > n \geq 1$ be integers. A smooth map from an m-dimensional smooth manifold with no boundary into an n-dimensional smooth manifold with no boundary is said to be a *fold* map if at each singular point p, the map is represented as $(x_1, \cdots, x_m) \mapsto (x_1, \cdots, x_{n-1}, \sum_{k=n}^{m-i} {x_k}^2 - \sum_{k=m-i+1}^{m} {x_k}^2)$ for suitable coordinates and an integer $0 \leq i(p) \leq \frac{m-n+1}{2}$. For the singular point p, $i(p)$ is unique and called the *index* of p. The set consisting of all singular points of a fixed index of the map is a closed submanifold of dimension $n-1$ with no boundary of the m-dimensional manifold. The restriction map to the singular set is an immersion. A *special generic* map, which is explained before, is defined as a fold map such that the index of a singular point is always 0.

Explicit construction of fold maps on explicit manifolds are fundamental, important and challenging: for example it also leads us to understand several important classes of manifolds such as 7-dimensional or higher dimensional closed and simply-connected manifolds more in geometric and constructive ways. Through the challenge, we have obtained the following theorem. For classical theory on 7-dimensional homotopy spheres, see [1] and [11] for example.

Theorem 1 ([2], [3] and so on.). (1) Every 7-dimensional homotopy oriented sphere M of 28 types admits a fold map $f : M \to \mathbb{R}^4$ such that $f|_{S(f)}$ is an embedding satisfying $f(S(f)) = \{x \in \mathbb{R}^4 \mid ||x|| = 1, 2, 3\}$, that the index of each singular point is 0 or 1, and the preimage of a regular value in each

connected component of f is, empty, diffeomorphic to S^3, diffeomorphic to $S^3 \sqcup S^3$ and diffeomorphic to $S^3 \sqcup S^3 \sqcup S^3$, respectively.

(2) 7-dimensional homotopy sphere M admits a fold map $f : M \to \mathbb{R}^4$ such that $f|_{S(f)}$ is an embedding and that $f(S(f)) = \{x \in \mathbb{R}^4 \mid ||x|| = 1\}$ if and only if M is a standard sphere. A 7-dimensional homotopy sphere M admits a fold map $f : M \to \mathbb{R}^4$ such that $f|_{S(f)}$ is an embedding and that the index of each singular point is 0 or 1 as before and in addition the following properties hold if and only if M is the total space of a linear bundle whose fiber is S^3 over S^4 (7-dimensional oriented homotopy spheres of 16 types are represented in this way).
   (a) $f(S(f)) = \{x \in \mathbb{R}^4 \mid ||x|| = 1, 2\}$.
   (b) For any connected component $C \subset f(S(f))$ and a small closed tubular neighborhood $N(C)$, the bundle given by the projection represented as the composition of $f|_{f^{-1}(N(C))}$ with the canonical projection to C is trivial.
   (c) The preimage of a regular value in each connected component is, empty, diffeomorphic to S^3, and diffeomorphic to $S^3 \sqcup S^3$, respectively.

As special generic maps, fold maps of a suitable class of maps considered there affect the differentiable structures of the homotopy spheres. Note that fold maps here are *round* fold maps: a round fold map is a fold map such that the restriction to the singular set is an embedding and that the singular value set is concentric spheres. Note also that fold maps here have been obtained as specific cases of fold maps on manifolds represented as connected sums of total spaces of bundles over the standard sphere of a fixed dimension whose fibers are standard spheres in [2], [3] and [6] for example. Related to this fact, manifolds represented as connected sums of manifolds represented as products of standard spheres admit special generic maps into suitable Euclidean spaces whose dimensions are smaller in considerable cases: for example a manifold represented as a connected sum of finite copies of $S^{n-1} \times S^{m-n+1}$ admits a special generic map into $\mathbb{R}^n$. We can construct the map so that the restriction to the singular set is an embedding and that the image is a manifold represented as a boundary connected sum of finite copies of $S^{n-1} \times I$ where I denotes the closed interval.

We present the following recent results, presenting explicit fold maps on infinitely many 7-dimensional closed and simply-connected manifolds. Their integral cohomology rings are mutually non-isomorphic. These manifolds are not represented as connected sums of manifolds represented as products of standard spheres as before or we cannot know whether they are represented so in general.

A *crossing* of an immersion is the point in the target such that the preimage has at least two points.

Theorem 2 ([7] and [8]). Let $\{G_j\}_{j=0}^7$ be a sequence of free and finitely generated commutative groups such that $G_0 = G_7 = \mathbb{Z}$, that G_j is zero for $j = 1, 6$, that G_j and G_{7-j} are mutually isomorphic for $j = 2, 3$ and that the rank of G_2 is smaller than or equal to that of G_4.

(1) In this situation, there exist a closed, simply-connected and spin manifold M of dimension 7 such that the homology group is free and that $H_j(M; \mathbb{Z})$ is isomorphic to G_j and a fold map $f : M \to \mathbb{R}^4$ such that $f|_{S(f)}$ is embedding, that the index of each singular point is always 0 or 1, and that

for each connected component of the regular value set of f, the preimage of a regular value in each connected component is, empty, diffeomorphic to S^3, or diffeomorphic to $S^3 \sqcup S^3$. Furthermore, if G_2 and G_4 are non-trivial groups, then there exist infinitely many closed and simply-connected manifolds M_λ of dimension 7 admitting the fold maps as before such that $H_j(M_\lambda; \mathbb{Z})$ is isomorphic to G_j and that the cohomology rings of M_{λ_1} and M_{λ_2} are not isomorphic for distinct $\lambda_1, \lambda_2 \in \Lambda$.

(2) In this situation, there exist a closed, simply-connected and spin manifold M of dimension 7 such that the homology group is free and that $H_j(M; \mathbb{Z})$ is isomorphic to G_j and a fold map $f : M \to \mathbb{R}^4$ such that $f|_{S(f)}$ is an immersion which may have crossings, that the index of each singular point is always 0 or 1, and that for each connected component of the regular value set of f, the preimage of a regular value in each connected component is, empty, diffeomorphic to S^3, diffeomorphic to $S^3 \sqcup S^3$, or diffeomorphic to $S^3 \sqcup S^3 \sqcup S^3$. Furthermore, if G_2 and G_4 are non-trivial groups, then there exist infinitely many closed and simply-connected manifolds M_λ of dimension 7 admitting the fold maps as before such that $H_j(M_\lambda; \mathbb{Z})$ is isomorphic to G_j and that the cohomology rings of M_{λ_1} and M_{λ_2} are not isomorphic for distinct $\lambda_1, \lambda_2 \in \Lambda$.

This result is seen as one capturing the cohomology rings of 7-dimensional closed and simply-connected manifolds of suitable families via explicit fold maps on them into $\mathbb{R}^4$.

The following theorem is the main theorem.

Main Theorem. There exists an infinite family $\{M_k\}_{k \in \mathbb{Z}}$ of closed, simply-connected and spin (oriented) manifolds whose integral cohomology rings are isomorphic to that of $\mathbb{C}P^2 \times S^3$ such that the 1st-Pontryagin class of M_k is $4k$ times a generator of $H^4(M_k; \mathbb{Z}) \cong \mathbb{Z}$ and these manifolds admit round fold maps $\{f_k : M_k \to \mathbb{R}^4\}$. Furthermore, every 7-dimensional, closed, simply-connected and spin manifold whose integral cohomology ring is isomorphic to that of $\mathbb{C}P^2 \times S^3$ admits a round fold map into $\mathbb{R}^4$.

1.3. **The content of the paper and acknowledgement.** The present paper consists of two sections and the remaining section is devoted to the proof of the main theorem. We construct a round fold map on a 7-dimensional, closed and simply-connected manifold and see that this is cohomologically $\mathbb{C}P^2 \times S^3$. We also present related results. We do not need to understand technique of construction of explicit fold maps such as arguments in [7], [8], and so on well.

2. The main theorem and related results.

We will prove the main theorem. We introduce fundamental terminologies.

For a finitely generated and free commutative group G, let $a \in G$ be a non-zero element which we cannot represent as $a = ra'$ for any pair $(r, a') \in (\mathbb{Z} - \{-1, 0, 1\}) \times G$ and let G be the internal direct sum of the group generated by the one element set $\{a\}$ and a subgroup of G. A homomorphism $a^* : G \to \mathbb{Z}$ is said to be the *dual* of a if $a^*(a) = 1$ and for any subgroup G' such that G is the internal direct sum of the group generated by the one element set $\{a\}$ and G', $a^*(G') = \{0\}$.

For a closed and orientable manifold (which is oriented), consider an integral homology class of degree k. It is said to be *represented* by a closed and connected

submanifold of dimension k with no boundary if the class is equal to the value of the homomorphism canonically induced from the inclusion at the *fundamental class* of the submanifold with an orientation: the *fundamental class* is a generator of the integral cohomology of degree k (of the k-dimensional closed manifold).

2.1. **The 3-dimensional complex projective space and its structure.** The 3-dimensional complex projective space $\mathbb{C}P^3$ is regarded as the total space of a linear bundle over S^4 whose fiber is S^2. It is a spin manifold. The 1st Pontryagin class of the projective space is 4 times a generator of the 4-the integral cohomology group. For this, see [10] for example. For classical systematic theory of Pontryagin classes see [13]. The following theorem presents several classical and important properties.

Theorem 3. For the projection $\pi_{\mathbb{C}P^3} : \mathbb{C}P^3 \to S^4$ of the linear bundle over S^4 whose fiber is S^2, the following properties hold.

(1) There exists a complex projective plane ${\mathbb{C}P^2}_{\mathbb{C}P^3} \subset \mathbb{C}P^3$ being a complex submanifold. Furthermore, a generator of $H_4(\mathbb{C}P^3;\mathbb{Z})$, isomorphic to $\mathbb{Z}$, is represented by the submanifold.
(2) $\pi_{\mathbb{C}P^3}|_{{\mathbb{C}P^2}_{\mathbb{C}P^3}}$ is a covering on the preimage of a smoothly embedded 4-dimensional standard closed disc in S^4.
(3) There exists a complex projective line ${\mathbb{C}P^1}_{\mathbb{C}P^2} \subset {\mathbb{C}P^2}_{\mathbb{C}P^3}$ being a complex submanifold and representing the generator of $H_2(\mathbb{C}P^3;\mathbb{Z})$, isomorphic to $\mathbb{Z}$. The complex projective line can be taken as a fiber of the bundle.
(4) The square of the dual of the integral homology class represented by ${\mathbb{C}P^1}_{\mathbb{C}P^2}$ with a suitable orientation is the dual of the integral homology class represented by ${\mathbb{C}P^2}_{\mathbb{C}P^3}$ with a suitable orientation. The product of the dual of the integral homology class represented by ${\mathbb{C}P^1}_{\mathbb{C}P^2}$ with a suitable orientation and the dual of the integral homology class represented by ${\mathbb{C}P^2}_{\mathbb{C}P^3}$ with a suitable orientation is a generator of $H^6(\mathbb{C}P^3;\mathbb{Z})$, isomorphic to $\mathbb{Z}$.

2.2. **A proof of the main theorem.** We consider a submersion obtained by composing the canonical projection from $\mathbb{C}P^3 \times S^1$ to $\mathbb{C}P^3$ with $\pi_{\mathbb{C}P^3} : \mathbb{C}P^3 \to S^4$ in Theorem 3. We have a 7-dimensional, closed and simply-connected manifold M and a fold map $f_{S^4} : M \to S^4$ by exchanging a restriction to the preimage of a smoothly embedded 4-dimensional standard disc D so that the following properties hold. Let $D' \subset S^4$ be a smoothly embedded 4-dimensional standard disc containing the disc D in the interior.

(1) f_{S^4} is a fold map such that the singular set $S(f_{S^4})$ is diffeomorphic to S^3, that $f_{S^4}|_{S(f_{S^4})}$ is an embedding and that $f_{S^4}(S(f_{S^4})) = \partial D'$.
(2) The preimage of a regular value of each connected component of $S^4 - f_{S^4}(S(f_{S^4}))$ is S^3 and $S^2 \times S^1$, respectively.

Note that in the second property, (a copy of) S^3 is regarded as a manifold obtained by a so-called handle attachment to (a copy of) $S^2 \times S^1$. In other words, we attach a *3-handle* $D^3 \times D^1$ along $S^2 \times \{*\} \subset S^2 \times S^1$, respecting the structure of a Morse function. See [12] for example for related theory. The following show topological properties of M obtained by investigating the two properties of the map before.

(1) There exists an oriented complex projective plane ${\mathbb{C}P^2}_M \subset M$ being a submanifold such that $f_{S^4}|_{{\mathbb{C}P^2}_M}$ is a covering on the preimage of the disc D in S^4.

(2) The integral cohomology ring of M is isomorphic to that of $\mathbb{C}P^2 \times S^3$.
(3) The integral homology class represented by the submanifold we can regard $S^2 \times \{*\} \subset S^2 \times S^1$ of the preimage of a regular value diffeomorphic to $S^2 \times S^1$ is a generator of $H_2(M;\mathbb{Z}) \cong \mathbb{Z}$. The integral homology classes represented by the preimages of regular values are a generator of $H_3(M;\mathbb{Z}) \cong \mathbb{Z}$. The square of the dual of the integral homology class represented by $S^2 \times \{*\}$ before with a suitable orientation is the dual of the integral homology class represented by ${\mathbb{C}P^2}_M \subset M$.
(4) M is spin.

We take the union of two smoothly and disjointly embedded 4-dimensional standard closed discs $D_1 := D$ and D_2 so that they are mutually in distinct connected components of the regular value set of f_{S^4}. By composing the restriction of f_{S^4} to the preimage ${f_{S^4}}^{-1}(D_1 \sqcup D_2)$ with a natural 2-fold covering over a 4-dimensional standard closed disc D_0, we have a trivial bundle over D_0 whose fiber is diffeomorphic to $S^3 \sqcup S^2 \times S^1$. We consider the projection and embed the base space as the space $\{x \in \mathbb{R}^4 \mid |x| \leq r_0\}$ for a positive integer $r_0 > 0$. Let $\bar{f_{r_0}}$ denote the resulting submersion. The restriction of f_{S^4} to ${f_{S^4}}^{-1}(S^4 - \mathrm{Int}(D_1 \sqcup D_2))$ is regarded as a product map of a Morse function with exactly one singular point and the identity map on id_{S^3}. On the domain of this, we can construct a product map of a Morse function $\bar{f}$ and the identity map on id_{S^3} such that the Morse function $\bar{f}$ satisfies the following properties.

(1) The function is a function on a manifold diffeomorphic to a manifold obtained by removing the interior of a 4-dimensional standard closed disc smoothly embedded in the interior of $S^2 \times D^2$.
(2) The minimum is r_0.
(3) The preimage of the minimum r_0 of the function coincides with the boundary and contains no singular point.
(4) The function has exactly three singular points and at distinct singular points, the values are distinct.

We can set the target of the function $\bar{f}$ as the half-closed interval $[r_0, +\infty) \subset \mathbb{R}$. Thanks to the structures of manifolds and maps, we can glue the maps $\bar{f_{r_0}}$ and the product map of $\bar{f}$ and the identity map on the boundary of $\{x \in \mathbb{R}^4 \mid |x| \leq r_0\}$ to obtain a global fold map on the original manifold M into $\mathbb{R}^4$.

Furthermore, the topological properties of M and $f_{S^4} : M \to S^4$ presented before together with fundamental propositions on 1st Pontryagin classes enable us to construct a similar fold map on a manifold whose integral cohomology ring is isomorphic to that of M (, which is oriented suitably,) and whose 1st Pontryagin class is $4k$ times a generator of the 4-th integral homology group, isomorphic to $\mathbb{Z}$, for an arbitrary integer k. We need to change the way we glue the trivial bundle over D_0 whose fiber is diffeomorphic to S^3. For Pontryagin classes see [13] again and see also Theorems 3 and 4 of [7].

This yields the following main theorem. A *round* fold map is a fold map such that the restriction to the singular set is an embedding and that the singular value set is concentric. The class of round fold maps was first defined by the author in the 2010s and have been systematically studied. See [2], [3], [4], [5] and [6] for more detailed presentations.




FIGURE 1. The image and the preimages of regular values of a round fold map in Theorem 4: circles represent the singular value set and 3-dimensional spheres.

Theorem 4. There exist an infinite family $\{M_k\}_{k\in\mathbb{Z}}$ of closed, simply-connected and spin (oriented) manifolds whose integral cohomology rings are isomorphic to that of $\mathbb{C}P^2 \times S^3$ such that the 1st-Pontryagin class of M_k is $4k$ times a generator of $H^4(M_k;\mathbb{Z}) \cong \mathbb{Z}$ and a family of round fold maps $\{f_k : M_k \to \mathbb{R}^4\}$ satisfying the following properties.

(1) The singular set of each map consists of exactly three connected components. More precisely, we can construct the maps so that the restrictions to the singular sets are embeddings and that the images are $\{x \in \mathbb{R}^4 \mid ||x|| = 1, 2, 3\}$.
(2) The preimage of a regular value in each connected component of the regular value set in the image is diffeomorphic to S^3, $S^2 \times S^1$ and $S^3 \sqcup S^2 \times S^1$, respectively.

See also FIGURE 1.

Theorem 5 ([18] and related fundamental and classical results). (1) For an arbitrary 7-dimensional, closed, simply-connected and spin manifold X whose integral cohomology ring is isomorphic to that of $\mathbb{C}P^2 \times S^3$, consider an isomorphism ϕ of integral cohomology rings from the integral cohomology ring of $\mathbb{C}P^2 \times S^3$ onto that of X and set a as a generator of $H^2(\mathbb{C}P^2 \times S^3;\mathbb{Z}) \cong \mathbb{Z}$. In this situation, the 1st Pontryagin class of X is represented as $4k\phi(a)$ for some integer k. Moreover, the topology of a 7-dimensional closed, simply-connected and spin manifold whose integral cohomology ring is isomorphic to that of $\mathbb{C}P^2 \times S^3$ is determined by its 1st Pontryagin class.
(2) For any pair of mutually homeomorphic 7-dimensional, closed, simply-connected and spin manifolds whose integral cohomology rings are isomorphic to that of $\mathbb{C}P^2 \times S^3$, one of the two manifolds is represented as a connected sum of the other manifold and a suitable 7-dimensional homotopy sphere. Moreover, for a 7-dimensional, closed, simply-connected and spin manifold whose integral cohomology ring is isomorphic to the integral cohomology ring of $\mathbb{C}P^2 \times S^3$, consider a pair of manifolds each of which is represented as a connected sum of the given manifold and a homotopy sphere. They are diffeomorphic if and only if the homotopy spheres are diffeomorphic.

Corollary 1. A 7-dimensional, closed, simply-connected and spin manifold whose integral cohomology ring is isomorphic to that of $\mathbb{C}P^2 \times S^3$ always admits a round fold map into $\mathbb{R}^4$.

Proof. Theorem 5 implies that a 7-dimensional, closed, simply-connected and spin manifold whose integral cohomology ring is isomorphic to that of $\mathbb{C}P^2 \times S^3$ is always




FIGURE 2. Construction of a round fold map on the manifold represented as a connected sum of the two manifolds admitting fold maps in Corollary 1: for example manifolds represent preimages of regular values in the connected components of the regular value sets. More precisely, we attach maps presented in FIGURE 3 and FIGURE 4 to obtain the desired map.

represented as a connected sum of a manifold in Theorem 4 and a 7-dimensional homotopy sphere. [3] and also [2], [6], and so on, show construction of a new round fold map from a round fold map in Theorem 4 and a round fold map in Theorem 2. We take a 4-dimensional, standard and closed disc in the connected component of the regular value set of the former map in the center and remove the interior of a connected component, regarded as the total space of a trivial bundle over the disc whose fiber is S^3 of the preimage. We remove (the preimage of) the union of the interior of a small closed tubular neighborhood of the outermost connected component of the singular value set of the latter map and the complementary set of the image. We glue the two obtained maps in a suitable way to obtain a round fold map on the desired manifold. See also FIGURE 2, FIGURE 3 and FIGURE 4. This completes the proof. □

Remark 1. The author believes that we can prove the following fact: 7-dimensional manifolds of Theorems 3, 4 and 5 and Main Theorem do not admit special generic maps or fold maps into $\mathbb{R}^n$ (n=1,2,3,4,5,6) such that the following hold (as in Theorem 1, Theorem 2, and so on).

(1) The index of each singular point is 0 or 1.
(2) Preimages of regular values are disjoint unions of standard spheres.

3. Acknowledgement.

This work is supported by "The Sasakawa Scientific Research Grant" (2020-2002 : https://www.jss.or.jp/ikusei/sasakawa/). The author is also a member of and




FIGURE 3. The image of a round fold map in Theorem 4 and the image of the map obtained by removing a connected component of the preimage of the interior of a smoothly embedded 4-dimensional standard closed disc.




FIGURE 4. The image of a round fold map in Theorem 2 and the image of the map obtained by removing the union of the complementary set of the image and the interior of a small closed tubular neighborhood of the outermost connected component of the singular value set and the preimage.

supported by JSPS KAKENHI Grant Number JP17H06128 "Innovative research of geometric topology and singularities of differentiable mappings".

REFERENCES

[1] J. J. Eells and N. H. Kuiper, *An invariant for certain smooth manifolds*, Ann. Mat. Pura Appl. 60 (1962), 93–110.
[2] N. Kitazawa, *On round fold maps* (in Japanese), RIMS Kokyuroku Bessatsu B38 (2013), 45–59.
[3] N. Kitazawa, *On manifolds admitting fold maps with singular value sets of concentric spheres*, Doctoral Dissertation, Tokyo Institute of Technology (2014).
[4] N. Kitazawa, *Fold maps with singular value sets of concentric spheres*, Hokkaido Mathematical Journal Vol.43, No.3 (2014), 327–359.
[5] N. Kitazawa, *Constructions of round fold maps on smooth bundles*, Tokyo J. of Math. Volume 37, Number 2, 385–403, arxiv:1305.1708.

[6] N. Kitazawa, *Round fold maps and the topologies and the differentiable structures of manifolds admitting explicit ones*, submitted to a refereed journal, arXiv:1304.0618 (the title has changed).
[7] N. Kitazawa *Notes on explicit smooth maps on 7-dimensional manifolds into the 4-dimensional Euclidean space*, submitted to a refereed journal, arxiv:1911.11274.
[8] N. Kitazawa *Explicit smooth maps on 7-dimensional closed and simply-connected manifolds of new classes*, arxiv:2005.05281.
[9] M. Kreck, *On the classification of* 1*-connected* 7*-manifolds with torsion free second homology*, to appear in the Journal of Topology, arxiv:1805.02391.
[10] R. D. Little, *The Pontrjagin class of a homotopy complex projective space*, Topology and its Applications 34 (1990), 257–267, https://core.ac.uk/download/pdf/82665808.pdf.
[11] J. Milnor, *On manifolds homeomorphic to the* 7*-sphere*, Ann. of Math. (2) 64 (1956), 399–405.
[12] J. Milnor, *Lectures on the h-cobordism theorem*, Math. Notes, Princeton Univ. Press, Princeton, N.J. 1965.
[13] J. Milnor and J. Stasheff, *Characteristic classes*, Annals of Mathematics Studies, No. 76. Princeton, N. J; Princeton University Press (1974).
[14] O. Saeki, *Topology of special generic maps of manifolds into Euclidean spaces*, Topology Appl. 49 (1993), 265–293.
[15] O. Saeki, *Topology of special generic maps into* $\mathbb{R}^3$, Workshop on Real and Complex Singularities (Sao Carlos, 1992), Mat. Contemp. 5 (1993), 161–186.
[16] O. Saeki and K. Sakuma, *On special generic maps into* $\mathbb{R}^3$, Pacific J. Math. 184 (1998), 175–193.
[17] O. Saeki and K. Sakuma, *Special generic maps of* 4*-manifolds and compact complex analytic surfaces*, Math. Ann. 313, 617–633, 1999.
[18] X. Wang *On the classification of certain* 1*-connected* 7*-manifolds and related problems*, arXiv:1810.08474.
[19] D. J. Wrazidlo, *Standard special generic maps of homotopy spheres into Eucidean spaces*, Topology Appl. 234 (2018), 348–358, arxiv:1707.08646.

Institute of Mathematics for Industry, Kyushu University, 744 Motooka, Nishi-ku Fukuoka 819-0395, Japan, TEL (Office): +81-92-802-4402, FAX (Office): +81-92-802-4405,

Email address: n-kitazawa@imi.kyushu-u.ac.jp
Webpage: https://naokikitazawa.github.io/NaokiKitazawa.html